

Antioxidant Enzymes Activities, Semen Characteristics and Fertility of Local Cocks in Relationship with Glutathione Levels and Storage Period

S. Shamiah, Ragaa Abd El-Karim and Abeir Eshera

Animal Production Research Institute, Agricultural Research Center, Giza, Egypt

THIS STUDY was carried out to evaluate the effect of glutathione (GSH) supplementation at levels of 0.0 (control), 0.2, 0.4 and 0.6 mM to diluted semen stored for 0, 24, 48 and 72 hr at 5°C, on antioxidant enzymes activities in seminal plasma, semen characteristics and fertility rate (FR) of cocks Inshas strain. Supplementation of 0.6 mM GSH to the diluted cocks semen significantly increased ($P<0.05$) the concentration of alkaline phosphatase (ALP) (40.2 u/ml) compared to other treatments (33.0, 27.8 and 28.3 u/ml). Superoxide dismutase (SOD) concentration in seminal plasma was not significantly affected ($P<0.05$) by different levels of GSH, storage period or interaction between levels of GSH and storage period. Glutathione peroxidase (GSH-Px) concentration in seminal plasma was not affected by different levels of GSH ($P<0.05$). Total antioxidant capacity (TACO) in seminal plasma was significantly affected ($P<0.05$) by different levels of GSH, storage period or interaction between them. There was a positive correlation ($r=0.089$ and $r = 0.168$) between SOD concentration and (GSH-Px and TAOC) concentration, but negative correlation ($r=-0.119$) with ALP concentration. Positive insignificant correlation between (ALP, SOD and TACO) concentration in seminal plasma with the progressive motility, live sperm, curled tail and fertility, while positive significant correlation between GSH-Px concentration with progressive motility, curled tail and fertility. It could be concluded that the addition of GSH to semen extender of cockerels especially at levels 0.2 mM improved the antioxidant enzymes activities, which play a vital role in lowering the deleterious effect of reactive oxygen species (ROS) on semen quality.

Keyword: Semen, Glutathione, *In vitro* storage, Antioxidant enzymes activity, Chicken

Introduction

The increasing use of artificial insemination (AI) in poultry industry emphasizes the need for good quality sperm distribution. A proper evaluation of semen prior to AI or storage is very important to avoid losses in fertility. Because semen evaluation is extremely important for semen storage and AI programs, a rapid, economical, objective and fertility predictive method of semen evaluation would be beneficial for the poultry industry protocols. Determination of antioxidant enzymes concentration in semen has been practiced for a long time (Peters, 2008).

Lipids are a basic component of semen, contributing to the membrane structure of

spermatozoa, the metabolism of the sperm cells, and their ability to capacitate and fertilize the female gamete. In birds, the lipid composition of spermatozoa has an influence on fertility (Ansah and Buckland, 1982). El-Saadany (2002) reported that increasing storage time of diluted semen from local chicken breed for 0, 24, 48 and 72 hr decreased significantly ($P<0.01$) the fertility rate (FR) by 5.12, 9.55 and 12.53%, respectively. Shamiah et al. (2017) reported that GSH supplementation in semen extender significantly ($P<0.05$) increased FR. Semen supplemented with 0.2 mM GSH (74.5%) showed higher FR than those supplemented with 0.6 mM GSH or free semen (59.9 and 59.2%, respectively).

Sperm cells have a high content of unsaturated

fatty acids in their membranes and they lack a significant cytoplasmic component containing antioxidants (Storey, 1997). Exposure of sperm to reactive oxygen species (ROS) resulted in lipid peroxidation (LPO), membrane breakdown, decreasing motility, abnormal morphology and a lowered capacity for oocytes penetration (Potts, et al., 1999). Deleterious effects of lipid peroxidation on poultry sperm include morphological defects, reduced motility, and poor fertilizing ability (Long and Kramer, 2003).

Several antioxidants such as glutathione (GSH), glutathione peroxidase (GSH-Px), catalase (CAT), and superoxide dismutase (SOD) act as a defense mechanism against the LPO and play a major role in maintaining sperm motility and viability (Bucak et al., 2008). One of the substances that is widely distributed in other types of cells is glutathione (GSH), where one of its functions is to protect cells against the destructive effect of reactive oxygen species (ROS, Meister, 1983). To counter the destructive effects of ROS, seminal plasma has antioxidant systems composed of SOD, CAT, GSH and GSH-Px (Aitken & Baker, 2004 and Gadea et al., 2004). However, sperm membrane has a large content of unsaturated fatty acids and lacks a significant cytoplasmic component containing antioxidants, and is particularly susceptible to LPO in the presence of ROS, leading to impaired cell function and decreased sperm motility (Bucak and Tekin, 2007).

Glutathione is a tripeptide L- γ -glutamyl-L-cysteinylglycine and ubiquitously distributed in organisms and plays a very important role in intracellular defence system against oxidative stress (Gadea et al., 2004). Glutathione could protect sperm due to its strong antioxidant activities by inhibiting the lipid peroxidation of red deer sperm during freezing-thawing processes. Glutathione was reported to be used as a cryoprotectant for the sperm of mouse, canine, boar, ovine, bull and red deer (Sławeta & Laskowska, 1987, Takeo & Nakagata 2011, Perez et al., 2012, Anel-López et al., 2015 and Lucio et al., 2016). Recently, Wang and Dong (2017) reported that glutathione addition to the extender improved sperm quality, including sperm motility, acrosomal integrity, plasma-membrane integrity and sperm morphology.

So, the aim of the present study was to evaluate

of the effect of GSH supplementation at levels of 0, 0.2, 0.4 and 0.6 mM to diluted semen stored for 0, 24, 48 and 72 hr at 5°C, on antioxidant enzymes activities and its relationship with sperm characteristics and fertility rate of Inshas strain of chicken.

Material and Methods

This study was carried out at Sakha Experimental Research Station, affiliated to the Animal Production Research Institute, Agricultural Research Center, Ministry of Agriculture.

Experimental chickens

Ten mature cockerels of local chicken (Inshas strain), 9-10 months of age and 2.30 -2.53 kg live body weight were used in this experiment. All cocks were kept in an individual cages (60x50x75 cm), provided with laying mash diet containing 15.5% crude protein and 2700 kcl ME per kg diet. Water was available all daytime.

Semen collection

Semen was collected in the morning from each cockerel once weekly by an abdominal massage method according to Lake and Stewart (1978) for five consecutive weeks as a semen collection period. Care was taken to avoid any contamination of semen with cloacae products and yellow and abnormal ejaculates were systematically discarded.

Only ejaculates with mass motility $\geq 70\%$ on the day of collection were pooled on each collection day. Semen was evaluated for sperm cell concentration using a Neubauer hemacytometer, then semen was extended by the modified Beltsville Poultry Semen Extender (mBPSE) at a rate of 1 semen: 2 extender and divided into four parts, the 1st part was extended with control extender without any supplementation (T1). However, the 2nd, 3rd and 4th parts were supplemented with 0.2 (T2), 0.4 (T3) and 0.6 (T4) mM GSH (C10H17N3O6S), respectively, (S.d fine-Chem. Limited, India).

Semen extender supplemented with different GSH levels was stored in refrigerator at 5°C and evaluated after 0, 24, 48 and 72 hr. Semen evaluation included PSM, SL, SA, acrosome status, and Hypo Osmotic Swelling Test (HOST) of spermatozoa.

Semen evaluation

The semen quality evaluation was performed for all treatment samples at different *in vitro* storage periods. The progressive motility percentage was assessed according to Tabatabaei et al. (2009). Sperm livability percentage was determined using eosin/nigrosin stain according to Lukaszewicz et al. (2008). Sperm abnormalities acrosome integrity was determined according to Al-Daraji (2001). Sperm membrane integrity was assessed by Hypo-osmotic swelling test (HOST) as described by Lagares et al. (2000). From each treatment at different storage time, 0.3 milliliters were taken and centrifuged at 3000 g for 10 min at 4°C and the supernatant was frozen and stored at -20°C at least 24 hr until analysis.

Antioxidant profiles assays

Semen TAOC, SOD, GSH-Px and ALP activities were measured by commercially available diagnostic kits, (MDSS GmbH Schiffgraben 41, 30175 Hannover, Germany).

Fertility trial

Artificial insemination was performed using 1.0 ml syringes for the deposition of the semen as described by Sadanand et al. (2004) to assess fertility rate. Hens from the same local bred were divided into 16 groups (7 hens for each treatment), including semen supplemented with 4 GSH level at 4 storage periods). A Total of 256 eggs, 16 from each group was collected after 1 day post-insemination for a week and incubated to evaluate the fertility rate (FR). Eggs were candled to identify the fertile eggs according to the method of Islam et al. (2002) as the following: FR = Fertilized eggs/ incubated eggs x100.

Statistical analyses

Data obtained were statistically analyzed as a factorial design (4 GSH levels x 4 storage periods) by analysis of variance ANOVA (SAS, 2001). The significant differences were set at 0.05 probability level and tested using the Duncan's procedure (Duncan, 1955).

Results and Discussion

Semen diluted with 0.2 mM GSH showed higher percentages of progressive motility, live sperm and curled tail, and lower percentages of abnormal sperm and acrosomal damage as compared with other groups ($P<0.05$). Semen without GSH (control) showed the poorest

quality (Fig. 1). These results clearly indicate that addition of GSH to semen extender at a level of 0.2 mM had impact on all sperm characteristics studied, regardless storage period. Also, addition of GSH to semen extender at a level of 0.2 mM improved fertility rate.

Ansari et al. (2014) observed that the addition of 0.5mM glutathione in semen extender improved the motility, plasma membrane integrity, viability and acrosomal integrity of Sahiwal bull spermatozoa. These results are in agreement with Wang and Dong (2017) who reported that glutathione addition to the extender improved sperm quality, including sperm motility, acrosomal integrity, plasma-membrane integrity and sperm morphology. Similarly, Gadea et al. (2005) reported beneficial effect of GSH in cryopreserved boar spermatozoa in terms of acrosomal intactness and percentage of non-capacitated spermatozoa, corroborated the findings of this study. Post translational modification in protein-thiols may mediate reduction in intracellular GSH and thus this is replenished by the supplementation of GSH into the extender (Gadea et al., 2004). In contrast, several studies also reported non-significant effects of GSH supplementation for protecting the sperms from cryocapacitation like changes (Foote et al., 2002, Sariozkan et al., 2009 and Tuncer et al., 2010).

In turkey, Long and Conn (2012) found that addition of phosphatidylcholine as antioxidant to the semen reduced the harmful effects of lipid peroxidation during semen storage at 4°C. Also, administration of vitamins C, E, catalase (Amini et al., 2015a & b) and oleic acid (Eslami et al., 2016) to the rooster semen decreased the levels of malondialdehyde in seminal plasma and post-thawed of sperm.

Results presented in Fig. 1 show that overall mean of fertility rate (FR) of eggs after performing the artificial insemination of hens was affected by GSH level ($P<0.05$). GSH supplementation significantly ($P<0.05$) increased FR. Semen supplemented with 0.2 mM GSH (74.5%) showed higher FR than those supplemented with 6 mM GSH or free semen (59.9 and 59.2%, respectively). However, FR of semen extended with 0.4 mM GSH did not differ significantly from other GSH levels or from the control semen. It is known that semen quality and fertility reduced by advancing storage period (Shamiah et al., 2017). Results of

the present study indicated that addition of GSH to semen extender improved fertility rate. Moreover, FR was higher for semen supplemented with

0.2 mM GSH as compared with other semen treatments, being the highest for fresh semen 0.2 and 0.4 mM GSH (74.45 and 68.36%) (Fig. 1).




Fig. 1. Sperm characteristics of cockerel and Fertility rate (%) as affected by glutathione supplementation to semen extender stored at 5°C

The alkaline phosphatase (ALP) activity in seminal plasma of cockerel's was not significantly affected ($P < 0.05$) by storage time, while the general trend decreased by increasing storage period (Table 1). Analysis of variance indicated that the overall mean of ALP enzyme activity was highly significantly ($P < 0.001$) increased with increasing level of GSH, the ALP activity was significantly ($P < 0.001$) higher (40.2 U/ml) with level 0.6 mM of GSH than other treatments (33.0, 27.8 and 28.3 U/ml). In the present study, the ALP activity insignificantly decreased by increasing storage period. In contrast, El-Saadany (2002) found that the ALP activity in seminal plasma in two local strains of cockerel's was increased significantly ($P < 0.05$) by increasing storage period. While El-Saadany (2002) found that the ALP activity increased significantly ($P < 0.05$) by increasing level of the antioxidant. Reduced GSH is synthesized in a controlled manner intra-cellularly in all mammalian cells and is involved in detoxification, antioxidant defense, and modulation of cell proliferation (Lu, 2009). It directly reacts with free radicals and catalyses the reduction of toxic H_2O_2 into water and hydro-peroxides (Bilodeau *et al.*, 2001). Both removal as well as dilution of seminal plasma

during semen freezing is reported to cause reduction in GSH content. This results in reduced GSH availability to spermatozoa as they fail to synthesize GSH by their own (Gadea *et al.*, 2011). Further, in a previous study it is also shown that supplementation of GSH in the extenders prior to freezing minimized the oxidative damage caused by free radical generation (Gadea *et al.*, 2005).

Effect of interaction between different levels of GSH and storage time on ALP concentration in diluted avian semen was not significant (Table 1). Mohan *et al.* (2013) found that no significant difference was noticed in ALP activity between fresh and preserved seminal plasma of guinea fowl. Also, Mahapatra *et al.* (1994) noted that ALP activity remained constant in fresh vs preserved semen in broiler breeders chicken.

As shown in Table 1, GSH-Px activity in seminal plasma was not significantly ($P < 0.05$) affected by different levels of GSH, while the nearest value of GSH-Px concentration with levels (0.2 and 0.4 mM, 150.5 and 150.2 μ g protein) was higher than (0.0 and 0.6 mM GSH, 144.2 and 144.9 mU/m). The overall mean of GSH-Px concentration in seminal plasma was significantly

($P < 0.05$) higher in time (0 hr) than in time (72hr), but did not differ from that in time (24 and 48hr). Also, no significant differences were observed among storage period (24, 48 and 72hr). Results of the present study are in the line with El-Saadany (2002) who found that GSH-Px activity increased significantly ($P < 0.05$) by advantages storage period. Previous studies indicated that, seminal plasma possesses free radical-trapping activity

and expresses SOD and GSH-Px activities (Surai et al., 1998). The free radical trapping activity of the seminal plasma was twice that of blood plasma and did not change after plasma boiling or storage during 24 hf at 20°C (Surai et al., 1998a). On the other hand, Thananurak et al. (2015) reported that membrane of avian spermatozoa is rich in polyunsaturated fatty acid (PUFAs) and can easily undergo LPO in the presence of ROS.

TABLE 1. Enzymatic activity in seminal plasma of cockerel as affected by glutathione supplementation to semen extender stored at 5°C

Item	Alkaline phosphatase	Glutathione peroxidase	Superoxide dismutase	Total antioxidant capacity
GSH (mM)				
0	28.3 ^b	144.2	159.3	0.97 ^c
0.2	27.8 ^b	150.5	162.2	1.32 ^a
0.4	33.0 ^b	150.2	161.4	1.15 ^b
0.6	40.2 ^a	144.9	158.4	1.08 ^{bc}
Storage period (h)				
0	36.4	163.0 ^a	165.3	1.04 ^{bc}
24	32.3	152.1 ^{ab}	167.5	1.37 ^a
48	30.8	145.8 ^{ab}	161.7	1.18 ^b
72	29.9	128.9 ^b	147.8	0.92 ^c
GSH × Storage period				
0×0	29.36	156.6	164.2	0.81
0×24	29.03	147.6	165.7	1.07
0×48	27.42	145.4	162.4	1.28
0×72	27.48	127.1	144.7	0.71
0.2×0	29.03	168.4	166.9	1.02
0.2×24	28.19	155.7	170.1	1.86
0.2×48	27.07	148.3	163.4	1.39
0.2×72	26.99	129.7	148.5	1.02
0.4×0	39.49	169.1	165.7	1.19
0.4×24	32.36	156.1	168.7	1.34
0.4×48	31.18	146.2	162.9	1.04
0.4×72	29.04	129.4	148.3	1.01
0.6×0	47.66	158.0	164.2	1.15
0.6×24	39.69	149.1	165.6	1.20
0.6×48	37.40	143.1	158.2	1.01
0.6×72	36.05	129.3	145.5	0.95
SEM	±4.31	±19.87	±16.56	±0.1
Significance				
GSH	***	NS	NS	***
Storage period	NS	NS	NS	***
GSH × Storage period	NS	NS	NS	**

a and b: Means denoted within the same column with different superscripts for each effect are significantly different.

NS= non significantly, **= significantly at $P < 0.01$ and ***= significantly at $P < 0.001$

The overall mean of SOD concentration in seminal plasma was not affected significantly ($P < 0.05$) by the different levels of GSH, storage period or interaction between them (Table 1). In general, SOD concentration in seminal plasma showed trend decreased insignificantly by advancing of storage period, while at levels of GSH, slightly increased up to 0.2 mM and slightly decreased up to 0.6 mM of GSH. The seminal plasma is well endowed with an array of antioxidants that act as free radical scavengers to protect spermatozoa against oxidative stress (Zini *et al.*, 2000, Agarwal *et al.*, 2004 and Sanocka & Kurpisza, 2004). Seminal plasma contains a number of enzymatic antioxidants such as SOD and CAT (Peeker *et al.*, 1997, Mruk *et al.*, 2002 and Calamera *et al.*, 2003). Moreover, Perumal *et al.* (2013) found that the concentration of SOD and CAT was higher in liquid storage (5°C) within semen treated with antioxidant (taurine) and say the SOD activity of sperm samples is a good predictor of their survival time. But normally, seminal plasma is a potent source of this antioxidant, SOD (Kobayashi *et al.*, 1991). Superoxide dismutase is an antioxidant enzyme found in avian and mammalian semen (Froman and Thurston, 1998, Kalaiselvi and Panneerselvam, 1998, Surai *et al.*, 1998a). Its purpose is to scavenge for biological oxidants like ROS and protect cells from peroxidative damage (Froman and Thurston, 1998). Superoxide dismutase scavenges both extracellular and intracellular superoxide anion and prevents lipid peroxidation of the plasma membrane. SOD spontaneously dismutase (O_2^-) anion to form O_2 and H_2O_2 (De Lamirande and Gagnon 1995). In addition, it contains a variety of nonenzymatic antioxidants such as vitamin C, vitamin E, pyruvate, glutathione, and carnitine (Meucci *et al.*, 2003).

In Table 1, the overall mean of TACO concentration in seminal plasma was significantly affected ($P < 0.001$) by different levels of GSH, storage period or interaction between them. The overall mean of TACO concentration in seminal plasma was significantly ($P < 0.001$) highest in time (24hr) than in other times. Also, TACO concentration was significantly ($P < 0.001$) higher in time (48hr) than in time (72 h). But did not differ among storage period (0.0, 48 and 72 hr). The overall mean of TACO concentration in seminal plasma was significantly ($P < 0.001$) highest with level (0.2mM) of GSH than in other treatments. Also, TACO concentration was significantly

($P < 0.05$) higher with level (0.4mM) than with level (0.0 mM), but did not differ among levels of GSH (0.4, 0.6 and 0.0 mM) (Table 1). The interaction between levels of GSH and storage period was affected significantly ($P < 0.01$), the highest concentration of TACO (1.86 mM/ml) with level of GSH (0.2 mM) at time 24 h and the lowest concentration (0.71 mM/ml) with level of GSH (0.0 mM) at time 72 h (Table 1).

Free radicals or ROS are deleterious to cell membranes. Exposure of cell membranes to ROS induces lipid peroxidation causing membrane breakdown and loss of function (Halliwell & Gutteridge, 1984 and Thananurak *et al.*, 2015). Lipid peroxidation results when intracellular production of ROS overcomes the antioxidant defense mechanisms utilized by cells including sperm (Alvarez *et al.*, 1987, Alvarez and Storey, 1989), and an immediate accumulation of lipid peroxides occurs in the plasma membrane. Unlike somatic cells, which depend on cytoplasmic enzymes such as SOD and GSH-Px to defend against peroxidation, spermatozoa release most of their cytoplasm immediately prior to sperm motion, and as a result, lose their enzymatic protection (Wang *et al.*, 1997). Avian sperm cell membranes have a much greater concentration of polyunsaturated fatty acids than mammalian sperm cells (Cerolini *et al.*, 1997) and are therefore more susceptible to lipid peroxidation during *in vitro* handling and storage of sperm, which is the primary cause of fertility dysfunction (Cecil and Bakst, 1993). Thus, determining the level of sperm lipid peroxidation can be utilized as a biochemical index of semen quality (Jones *et al.*, 1978 and Alvarez *et al.*, 1987).

Table (2) shows slightly positive correlation ($r = 0.089$ and $r = 0.168$) between SOD concentration and (GSHPx and TAOC) concentration, but negative correlation ($r = -0.119$) with ALP concentration in seminal plasma of corresponding samples. Positive correlation significantly were found between TAOC concentration and (GSHPx and ALP), Also between concentration of GSHPx and ALP (Table 2). The total antioxidant capacity of seminal plasma represents the sum of the potential anti-ROS enzymes, such as GSH-PX. (Mahfouz, *et al.*, 2009) Furthermore, Contri *et al.*, (2011) reported a positive correlation between sperm parameters and total antioxidant capacity in seminal plasma.

TABLE 2. Correlation between some of antioxidants enzymes concentration in cocks seminal plasma.

Item	Alkaline phosphatase (ALP)	Glutathione peroxidase (GSHpx)	Total antioxidant capacity (TACO)
Glutathione peroxidase (GSHpx)	0.362*	1	
Total antioxidant capacity (TACO)	0.498**	0.673**	1
Superoxide dismutase (SOD)	-0.119	0.089	0.168

The positive correlation between ALP, SOD and TACO concentration in seminal plasma with the progressive motility, live sperm, curled tail and fertility were not significant, while positive significant correlation between GSH-Px concentration with progressive motility, curled tail

and fertility (Table 3). In rooster, supplementation of oleic to semen extender caused an improve in total antioxidant activity concentration of seminal plasma and spermatozoa stored at 4°C for at 24 h and 48h compared with control group (Eslami et al., 2016).

TABLE 3. Correlation between some of antioxidant enzymes concentration in cocks seminal plasma and semen characteristics.

Parameter	Alkaline phosphatase (ALP)	Glutathione peroxidase (GSHpx)	Total antioxidant capacity (TACO)	Superoxide Dismutase (SOD)
Progressive motility	0.166	0.298*	0.117	0.106
Live sperm	0.128	0.260	0.127	0.268
Abnormal sperm	-0.116	-0.22	-0.041	-0.346*
Acrosomal damage	-0.333*	-0.524**	-0.394**	-0.283
Curled tail	0.226	0.314*	0.159	0.142
Fertility	0.178	0.357*	0.242	0.27

Slightly negative correlation between (ALP, GSHPx and TAOC) concentration in seminal plasma and abnormal sperm was insignificant, but moderate negative correlation between (ALP, GSHPx and TAOC) concentration and acrosomal damage was significant also, SOD concentration with abnormal sperm (Table 3). Mammalian sperm is equipped with a defense system that comprises of both enzymatic and non-enzymatic antioxidants which offer protection against the ROS. But this indigenous defensive mechanism of the sperm is reported to be insufficient to cope with the oxidative stress (Nichi et al. 2006). Moreover, antioxidant levels of semen decrease as a result of cryopreservation (Bilodeau et al. 2000). However, addition of antioxidants in the extender has been reported to improve the quality of preserved mammalian semen (Bilodeau et al. 2001).

Glutathione is a tripeptide naturally occurring in semen and providing intracellular defense to the sperm against the oxidative stress caused by an over-production of ROS during the freezing

and thawing process. Freeze-thawing of the semen causes a significant reduction in the glutathione content of the boar (Gadea et al. 2004) and bull semen (Bilodeau et al. 2000). Moreover, supplementation of glutathione in the extender has been shown to provide a protective role in maintaining the quality of bovine (Gadea et al. 2007) and buck semen (Sinha et al. 1996).

Shamiah et al., (2017) reported that the semen supplemented with GSH was characterized by moderate resistance morphological defects, increases motility and livability of spermatozoa in preserved semen as well as fertility rates. Supplementation of GSH at a level of 0.2 mM to extender of cockerel semen stored at 5°C for up to 72 h is an appropriate level to improve the fertilizing ability of chicken spermatozoa. Also, in buffalo, El-Kon and Darwish (2011) indicating that addition of 0.50-1.00 mM of GSH to semen diluent improved quality of liquid semen stored up to 120 hours in terms of reducing DNA damage and improving the fertility.

Hydrogen peroxide is the primary toxic ROS for human spermatozoa that its high concentration induces lipid peroxidation and results in cell death. Therefore, the balance of the SOD and CAT activities in semen is important for maintaining sperm motility (Hsieh *et al.*, 2002), and complement, that there is a positive but no significantly correlation between SOD activity and sperm motility. They interpreted that higher SOD activity may scavenge the generation of ROS, which may lower the cytotoxicity to spermatozoa.

Conclusions

Results in the present study clearly indicated that the addition of GSH to semen extender of cockerels especially at levels 0.2 mM improved the antioxidant enzymes activity and these enzymes have a vital role in lowering the deleterious effect of reactive oxygen species (ROS) on semen quality.

References

- Agarwal A., Nallella K.P., Allamaneni S.S. and Said T.M., (2004) Role of antioxidants in treatment of male infertility: an overview of the literature. *Reprod. Biomed. Online*, **8**, 616-627.
- Aitken, R.J. and Baker, M.A. (2004) Oxidative stress and male reproductive biology. *Reprod. Fertil.*, **16**, 581-588.
- Al-Daraj, H.J. (2001) Effects of holding temperature and time on acrosomal abnormalities of fowl sperms. *Indian J. Anim. Sci.*, **71**, 32 – 34.
- Alvarez, J.G. and Storey, B.T. (1989) Role of glutathione peroxidase in protecting mammalian spermatozoa from loss of motility caused by spontaneous lipid peroxidation. *Gamete Res.*, **23**, 77-90.
- Alvarez, J. G., Touchstone, J.C., Blasco, L. and Storey, B.T. (1987) Spontaneous lipid peroxidation and production of hydrogen peroxide and superoxide in human spermatozoa. *J. Androl.*, **8**, 338-348.
- Amini, M.R., Kohram, H., Zare Shahaneh, A., Zhandi, M., Sharideh H. and Nabi, M.M. (2015a) The effects of different levels of vitamin E and vitamin C in modified Beltsville extender on rooster post-thawed sperm quality. *Cell Tissue Bank*. **16**, 587-592.
- Amini, M.R., Kohram, H., Zare Shahaneh, A., Zhandi, M., Sharideh, H. and Nabi, M.M. (2015b) The effects of different levels of catalase and superoxide dismutase in modified Beltsville extender on rooster post-thawed sperm quality. *Cryobiology*, **70**, 226-232.
- Anel-López L., Garcia-Alvarez O., Maroto-Morales A., Iniesta-Cuerda M., Ramón M., Soler A.J., Fernández-Santos M.R. and Garde, J.J. (2015) Reduced glutathione addition improves both the kinematics and physiological quality of post-thawed red deer sperm. *Anim Reprod Sci.* **162**, 73-79.
- Ansah, G.A. and Buckland, R.B. (1982) Genetic variation in fowl semen cholesterol and phospholipid levels and the relationships of these lipids with fertility of frozen-thawed and fresh semen, *Poult. Sci.* **61**, pp. 623-637.
- Ansari M. S., Rakha, B. Allah , Iqbal, R. and Akhter, S. (2014) Effect of Glutathione in Extender on the Freezability of Sahiwal Bull Spermatozoa. *Pakistan J. Zool.*, **46**, 17-21.
- Bilodeau J.F., Blanchette S., Gagnon C. and Sirad M.A. (2000) Levels of antioxidant defenses are decreased in bovine spermatozoa after a cycle of freezing and thawing. *Molecular Reproduction and Development*, **55**, 282-288.
- Bilodeau, J.F., Blanchette, S., Gagnon, C. and Siarrd, M.A. (2001) Thiols prevent H₂O₂ mediated loss of sperm motility in cryopreserved bull semen. *Theriogenology*, **56**, 275-286.
- Bucak M.N. and Tekin N. (2007) Protective effect of taurine, glutathione and trehalose on the liquid storage of ram semen. *Small Rumin. Res.* **73**, 103-108.
- Bucak, M.N., Atessahin, A. and Yüce, A. (2008) Effect of anti-oxidants and oxidative stress parameters on ram semen after the freeze-thawing process. *Small Rumin. Res.* **75**, 128-134.
- Calamera, J., Buffone, M., Ollero, M., Alvarez, J. and Doncel, G.F. (2003) Superoxide

- dismutase content and fatty acid composition in subsets of human spermatozoa from normozoospermic, asthenozoospermic, and polyzoospermic semen samples. *Mol. Reprod. Dev.* **66**, 422-430.
- Cecil, H.C. and Bakst, M.R. (1993) *In vitro* lipid peroxidation of turkey spermatozoa, *Poult. Sci.* pp. 1370-1378.
- Cerolini, S., Kelso, K. A., Noble, R. C., Speake, B. K., Pizzi, F. and Cavalchini, L. G. (1997) Relationship between spermatozoan lipid composition and fertility during aging of chickens. *Biol. Reprod.* **57**,976-980.
- Contri, A., De Amicis, I. and Molinari, A. (2011) Effect of dietary antioxidant supplementation on fresh semen quality in stallion. *Theriogenology*, **75**,1319-1326.
- Daryabari, H., Akhlaghi, A., Zamiri, M. J., Pirsaraei, Z.A., Mianji, G.R., Deldar, H. and Eghbalian, A.N. (2015) Oral administration of supplementary biotin differentially influences the fertility rate and oviductal expression of avidin and avidin-related protein-2 in low- and high fertility broiler line hens. *Poult. Sci.* **94**,289-295.
- De Lamirande E. and Gagnon, C. (1995) Impact of reactive oxygen species on spermatozoa: a balancing act between beneficial and detrimental effects, *Human Reproduction*, **10**, 15-21.
- Duncan, B. D. (1955) Multiple ranges and multiple F test. *Biometrics*, **11**, 1-42.
- El-kon, I.I. and Darwish, S.A. (2011) Effect of glutathione (GSH) on microscopic parameters and DNA integrity in Egyptian buffalo semen during liquid and frozen storage. *Journal of Reproduction and Infertility*, **3**, 32-40.
- El-Saadany, Amina, Sh. (2002) Use of antioxidants in storing local cockerels semen. *M.Sc. Thesis*, Alex. Univ., Egypt.
- Eslami, M., Ghaniei, A. and Mirzaei, R. H. (2016) Effect of the rooster semen enrichment with oleic acid on the quality of semen during chilled storage. *Poultry Sci.* **95**,1418-1424.
- Foote, R.H., Brockett, C.C. and Kaproth, M.T. (2002) Motility and fertility of bull sperm in whole milk extender containing antioxidants. *Anim. Reprod. Sci.* **71**, 13-23.
- Froman, D. P., and Thurston, R. J. (1998) Chicken and turkey spermatozoal superoxide dismutase: A comparative study. *Biol. Reprod.* **24**,193-200.
- Gadea J, Sellis, E., Marco, M.A., Coy, P., Matás, .C, Romar, R. and Ruiz, S. (2004) Decrease in glutathione content in boar sperm after cryopreservation: effect of the addition of reduced glutathione to the freezing and thawing extenders. *Theriogenology*, **62**,690-701.
- Gadea, J., Francisco, G.V., Carmen, M.S., Juan, C.G.N., Sebastia, N.C.N. and David, G. (2005) Cooling and freezing of boar spermatozoa: supplementation of the freezing media with reduced glutathione preserves sperm function. *J. Androl.* **26**, 396-404.
- Gadea J., Gumbao, D., Novas, S.C., Zquez, F.A.Z., Grullo, L.A. and Gardo, G.C. (2007) Supplementation of the dilution medium after thawing with reduced glutathione improves function and the in vitro fertilizing ability of frozen-thawed bull spermatozoa. *Andrology*, **7**, 1-10.
- Gadea, J., Molla, M., Selles, E., Marco, M.A., Garcia-Vazquez, F.A. and Gordan, J.C. (2011) Reduced glutathione content in human sperm is decreased after cryopreservation: effect of the addition of reduced glutathione to the freezing and thawing extenders. *Cryobiology*, **62**, 40-46.
- Halliwell, B. and Gutteridge, J. M. C. (1984) Lipid peroxidation, oxygen radicals, cell damage, and antioxidant therapy. *Lancet* **334**,1396-1397.
- Hsieh, Y.Y., Sun, Y.L., Chang, C.C., Lee, Y.S., Tsai, H.D. and Lin, C.S. (2002) Superoxide dismutase activities of spermatozoa and seminal plasma are not correlated with male infertility. *J. Clin. Lab Anal.* **16**,127-131.
- Islam, M. S., Howlider, M. A. R., Kabir, F. and Alam, J. (2002) Comparative assessment of fertility and hatchability of barred Plymouth

- rock, White Leghorn, Rhode Island Red and White Rock hens. *Int. J. Poult. Sci.* **1**, 85-90.
- Jones, R., Mann, T. and Sherins, R. J. (1978) Adverse effects of peroxidized lipid on human spermatozoa. *Proc. R. Soc. Lond.* **201**,413–417.
- Kalaiselvi, T. and Panneerselvam, C. (1998) Effect of L-carnitine on the status of lipid peroxidation and antioxidants in aging rats. *J. Nutr. Biochem.* **9**,575–581.
- Kobayashi, M., Kakizono, T. and Nagai, S. (1991) Astaxanthin production by a green alga, *Haematococcus pluvialis* accompanied with morphological changes in acetate media. *J. Ferment. Bioeng.* **71**, 335–339.
- Lagares, M.A., Petzoldt, R., Sieme, H. and Klug, E. (2000) Assessing equine sperm-membrane integrity. *Andrologia*, **32**,163-167.
- Lake, P.E. and Stewart, J.M. (1978) Artificial insemination in poultry. Bulletin, 213, Ministry of Agriculture Fisheries and Food, London.
- Long, J. A. and Conn, T. L. (2012) Use of phosphatidylcholine to improve the function of turkey semen stored at 4°C for 24 hours. *Poult. Sci.* **91**,1990–1996.
- Long, J.A. and Kramer, M. (2003) Effect of vitamin E on lipid peroxidation and fertility after artificial insemination with liquid-stored Turkey semen, *Poult. Sci.* **82**, pp. 1802–1807.
- Lu, S.C. (2009) Regulation of glutathione synthesis. *Mol. Asp. Med.* **30**, 42–59.
- Lucio C.F., Silva L.C.G., Regazzi, F.M., Angrimani, D.S., Nichi, M., Assumpção M.E., Vannucchi C.I. (2016) Effect of reduced glutathione (GSH) in canine sperm cryopreservation: in vitro and in vivo evaluation. *Cryobiology*, **72**,135–140.
- Lukaszewicz, E., Jersey, A., Partyka, A. and Siudzinska, A. (2008) Efficacy of evaluation of rooster sperm morphology using different staining methods. *Res. Vet. Sci.* **85**, 583-588.
- Mahapatra, P.S., Mohanty, B.P., Bisoi, P.C., Mishra, S.C., Nayak, N.R. and Mishra, M.S. (1994) Effect of storage on the physical and biochemical parameters of broiler semen. *Indian J. Poult. Sci.* **29**, 146-150.
- Mahfouz R., Sharma R. and Sharma D. (2009) Diagnostic value of the total antioxidant capacity in human seminal plasma. *Fertil. Steril.* **91**,805-811.
- Meister, A. (1983) Selective modification of glutathione metabolism. *Science*, **230**, 472 – 477.
- Mosenene, T. M. B. (2009) Characterization and cryopreservation of semen of four south african chicken breeds. M.Sc., Faculty of Natural and Agricultural Sciences, University of the Free State, Bloemfontein.
- Mruk D.D., Silvestrini B., Mo, M.Y. and Cheng, C.Y. (2002) Antioxidant superoxide dismutase – a review: its function, regulation in the testis, and role in male fertility. *Contraception*, **65**, 305-311.
- Nichi M., Bols, P.E.J., Zuge, R.M., Barnabe, V.M., Goovaerts, I.G.F., Barnabe, R.C. and Cortada, C.N.M. (2006) Seasonal variation in semen quality in *Bos indicus* and *Bos taurus* bulls raised under tropical conditions. *Theriogenology*, **66**, 822-828.
- Peeker, R., Abramsson, L. and Marklund, S.L. (1997) Superoxide dismutase isoenzymes in human seminal plasma and spermatozoa. *Mol. Hum. Reprod.* **3**,1061-1066.
- Perez EGA, Nichi M, Viana CHC, Silva R O., Dalmazzo A, Góes PA, Gurgel J R, Barnabe V, Barnabe R. (2012) Effect of glutathione on the function and oxidative status of ovine cryopreserved sperm. *Brazilian J. Vet. Res. Anim. Sci.*, **49**,262–268.
- Perumal, P., Kezhavituo Vupru and Rajkhowa, C. (2013) Effect of addition of taurine on the liquid storage (5°C) of mithun (*Bos frontalis*) semen. *Vet. Med. Int.* 1-7.
- Peters, S. O. (2008) Semen quality traits of seven strain of chickens raised in the humid tropics / Peters, S. O., Shoyebo, O. D., Ilori, B. M. et al. // *Int. J. Poult. Sci.* **7** (10), 949–953.
- Potts, R.J., Jefferies, T.M. and Notarianni, L.J.

- (1999) Antioxidant capacity of the epididymis. *Hum. Reprod.* **14**,2513–2516.
- Rosa, A. P., Scher, A., Sorbara, J. O. B., Boemo, L. S., Forgiarini, J. and Londero, A. (2012) Effects of canthaxanthin on the productive and reproductive performance of broiler breeders. *Poult. Sci.* **91**,660–666.
- Sadanand, D., Sontakke, G.U., Vedantam, S., Sanjiva, D. and Kholkute, S.S. (2004) Semen characteristics, cryopreservation and successful artificial insemination in the Blue rock pigeon (*Columba livia*). *Theriogenology*, **62**, 139-153.
- Sanocka D. and Kurpisz M. (2004) Reactive oxygen species and sperm cells. *Reprod. Biol. Endocrinol.* **2**,12.
- Sariozkan, S., Bucak, M.N., Tuncer, P.B., Ulutas, P.B., Bilgean, A. (2009) The influence of cysteine and taurine on microscopic-oxidative stress parameter and fertilizing ability of bull semen following cryopreservation. *Cryobiology*, **58**, 134–138.
- SAS, Institute. (2001) SAS User's Guide: Statistics version 6.12 ed. SAS Institute, Inc., Cary, NC., USA.
- Shamiah, Sh. M., Abd El-Karim, Ragaa E., Fouda, Sara F., Eshera, Abeir A.M. and El-Ratel, I.T. (2017) Effect of glutathione supplementation to semen extender on cockerel sperm characteristics in semen stored at 5°C for different periods. *Egyptian J. Anim. Prod.* **54**,17-22
- Sinha, M.P., Sinha, A.K., Singh, B.K. and Prasad, P.L. (1996) The effect of glutathione on the motility, enzyme leakage and fertility of frozen goat semen. *Theriogenology*, **41**, 237–243.
- Sławeta R. and Laskowska, T. (1987) The effect of glutathione on the motility and fertility of frozen bull sperm. *Anim. Reprod. Sci.* **13**, 249–253.
- Storey, B.T. (1997) Biochemistry of the induction and prevention of lipoperoxidative damage in human spermatozoa. *Mol. Hum. Reprod.* **3**, 203 – 214.
- Surai, P.F., Blesbois, E., Grasseau, I., Chalah, T., Brillard, J.P., Wishart, G.J., Cerolini S. and Sparks, N.H.C. (1998) Fatty acid composition, glutathione peroxidase and superoxide dismutase activity and total antioxidant activity of avian semen. *Comp. Biochem. Phys. B* **120**, 527–533.
- Tabatabaei, S., Batavani, R.A. and Talebi, A.R. (2009) Comparison of semen quality in indigenous and Ross broiler breeder roosters. *J. Anin. Vet. Adv.* **8**, 90–93.
- Takeo, T. and Nakagata, N. (2011) Reduced glutathione enhances fertility of frozen/thawed C57BL/6 mouse sperm after exposure to methyl-beta-cyclodextrin. *Biol. Reprod.* **85**,1066–1072.
- Thananurak, P., Sittikasamkit, C., Vongpralub, T. and Sakwivatkul, K. (2015) Effects of addition of reduced glutathione to thawing media on motility parameters, lipid peroxidation and fertility rate in frozen-thawed chicken spermatozoa. *Khon Kaen AGR. J.* **43** Suppl. 2, 98-102.
- Tuncer, P.B., Bucak, M.N., Serhat, B., Sariozkan, S., Yeni, D., Eken, A., Akalin, P.P., Kinet, H., Avdatek, F., Fidan, A.F. and Gundogan, M. (2010) The effect of cysteine and glutathione on sperm and oxidative stress parameter of post-thawed bull semen. *Cryobiology*, **61**, 303–307.
- Wang Y. and Dong, S. (2017) Glutathione in combination with trehalose has supplementary beneficial effects on cryopreserved red deer (*Cervus elaphus*) sperm. *Am. J. Reprod. Immunol.* **77**, e12610, DOI: 10.1111/aji.12610
- Wang, Y., Sharma, R.K. and Agarwal, A. (1997) Effect of cryopreservation and sperm concentration on lipid peroxidation in human semen. *Urology*, **50**,409–413.
- Zini, A., Garrels, K. and Phang, D. (2000) Antioxidant activity in the semen of fertile and infertile men. *Urology*, **55**, 922-926.

(Received : 13 / 8 / 2017)
 accepted : 18 / 9 / 2017)

تأثير مستويات الجلوتاثيون ومدة التخزين على النشاط الانزيمي لمضادات الاكسده وعلاقتها بخصائص السائل المنوي والخصوبة للديكة المحلية

شريف شامية ، رجاء عبدالكريم وعبير عشيبة

معهد بحوث الإنتاج الحيواني – مركز البحوث الزراعية – الجيزة - مصر.

تهدف هذه الدراسة الى تقييم تأثير اضافة الجلوتاثيون بمستويات صفر ٠,٢, ٠,٤, ٠,٦ و ٠,٦ ملل مول لمخفف السائل المنوي المخزن لمدة صفر ٠, ٢٤, ٤٨ و ٧٢ ساعة على درجة ٥٥°م على نشاط الانزيمات المضادة للاكسده في بلازما السائل المنوي وتحديد مدى ارتباطها مع خصائص الحيوانات المنوية ومعدل الخصوبة للديوك من سلالة انشاص. وقد اوضحت النتائج عدم وجود تأثير معنوي لانزيم الالكالين فوسفاتيز (ALP) بزيادة فترة التخزين على الرغم من أنه عند اضافة ٠,٦ ملل مول من الجلوتاثيون أدى إلى زيادة معنوية في تركيز انزيم الالكالين فوسفاتيز بمقدار (٤٠,٢ u/ml) مقارنة بالمعاملات الأخرى (٣٣,٠, ٢٧,٨, ٢٨,٣). كما لوحظ عدم تأثير نشاط إنزيم سوبراكسيد ديسيمتيز (SOD) في بلازما السائل المنوي عند اضافة اى مستوى من الجلوتاثيون او عند اى فترة تخزين وكذا عند التفاعل بينهما. لم يتأثر معنويا تركيز الجلوتاثيون بيروكسيديز في بلازما السائل المنوي باختلاف مستويات الجلوتاثيون المضافة، في حين اقتربت قيم GSH-Px بين المستويين ٠,٢ و ٠,٤ ملل مول جلوتاثيون حيث كانت ١٥٠,٥ و ١٥٠,٢ ميكرون /جرام بروتين أعلى من المستويين الكنترول (بدون جلوتاثيون) و ٠,٦ ملل مول جلوتاثيون حيث كانت ١٤٤,٢ و ١٤٤,٩ ميكرون /جرام بروتين. كما اوضحت الدراسة وجود تأثير معنوي لاضافة المستويات المختلفة من الجلوتاثيون مع طول فترة التخزين والتفاعل بينهما على المتوسط العام لتركيز TACO في بلازما السائل المنوي.

وجد ارتباط موجب ($r = 0,089$ ، $r = 0,168$) بين تركيز ALP، TAOC، GSH-Px، SOD، بينما وجد ارتباط سلبى ($r = -0,119$) مع تركيز ALP في بلازما السائل المنوي. كما وجد ارتباط معنوي موجب بين تركيز الإنزيمات المضادة للأكسدة في بلازما السائل المنوي و الحركة الجماعية و الحيوية و نسبة الحى، كما كان الارتباط ضعيفا بين تركيز الإنزيمات المضادة للأكسدة في البلازما المنوية مع الخصوبة، بينما وجد ارتباط سلبى كبير بين تركيز GSH-Px و الخصوبة

الخلاصة :

النتائج في هذه الدراسة تشير بوضوح الى ان اضافة الجلوتاثيون (GSH) الى مخفف السائل المنوي خاصة عند مستوى ٠,٢ ملل مول تحسن من نشاط الانزيمات المضادة للاكسده وان هذه الانزيمات لها دور فى تقليص الاثر الضار للشوارد (ROS) الحره على جودة السائل المنوي لديوك الدجاج.

